

Time variation of the fine structure constant and of the Higgs vacuum expectation value on cosmological time scales

M. E. Mosquera^{1,2} and O. Civitarese²

¹ Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque, (1900) La Plata, Argentina
e-mail: osvaldo.civitarese@fisica.unlp.edu.ar

² Department of Physics, University of La Plata c.c. 67 (1900), La Plata, Argentina

Received 16 November 2010 / Accepted 29 November 2010

ABSTRACT

Aims. We study the time variation of both the fine structure constant, α , and the Higgs vacuum expectation value, v , during Big Bang nucleosynthesis (BBN).

Methods. We calculate the primordial abundances of D, ^4He , and ^7Li by modifying Kawano's code, and by using observational data we set constraints on the joint variation of α and v . To perform the calculations we considered the dependence of the deuterium-binding energy, ϵ_D , upon v , obtained from the treatment of different proton-neutron interactions.

Results. Results are consistent with variation on v (even at the level of 6σ) and null variation of α (within 2σ) if the ^7Li data are used in the analysis. However, if data on these nuclei are not considered in the statistical analysis, we found null variation on both fundamental constants within 2σ .

Conclusions. We found that the best-fit values of the variation of v and α are sensible to the dependence of the deuterium binding energy upon the Higgs vacuum expectation value. We found non-null variation of v within 6σ if all the observational data are used in the analysis. If data on the primordial abundance of ^7Li are taken at face value, the discrepancy between BBN and WMAP estimates may be explained by allowing variations of v .

Key words. primordial nucleosynthesis – nuclear reactions – nucleosynthesis – abundances – cosmology: theory

1. Introduction

Several observations can establish limits to the variation of different fundamental constants, such as the atomic clocks (Bize et al. 2003; Fischer et al. 2004; Peik et al. 2004; Prestage et al. 1995; Sortais et al. 2000; Marion et al. 2003), the Oklo natural fission reactor (Damour & Dyson 1996; Fujii et al. 2000) and the analysis of the spectra of quasar absorption systems. These astronomical observations suggest a possible variation of the fine structure constant and the electron-to-proton mass ratio (Webb et al. 1999, 2001; Murphy et al. 2001a,b, 2003; Ivanchik et al. 2005; Reinhold et al. 2006; Tzanavaris et al. 2007). However, another analysis of similar astronomical data gives null variation of the fine structure constant (Martínez Fiorenzano et al. 2003; Quast et al. 2004; Bahcall et al. 2004; Srianand et al. 2004; King et al. 2008; Thompson et al. 2009; Malec et al. 2010).

The Big Bang nucleosynthesis (BBN) is a useful tool to study the time variation of fundamental constants, such as the fine structure constant, α , the Higgs vacuum expectation value, v , and the Planck mass, among others. Several theories that attempt to unify the four fundamental interactions, such as superstrings (Wu & Wang 1986; Barr & Mohapatra 1988; Maeda 1988; Damour & Polyakov 1994; Damour et al. 2002a,b), brane world (Youn 2001a,b; Brax et al. 2003; Palma et al. 2003) and Kaluza-Klein theories (Kaluza 1921; Klein 1926; Weinberg 1983; Gleiser & Taylor 1985; Overduin & Wesson 1997), allow fundamental constants to vary within cosmological time scales. The time variation of several fundamental constants was

studied by Campbell & Olive (1995), Bergström et al. (1999), Ichikawa & Kawasaki (2002), Nollett & Lopez (2002), Yoo & Scherrer (2003), Müller et al. (2004), Ichikawa & Kawasaki (2004), Cyburt et al. (2005), Landau et al. (2006), Coc et al. (2007), Chamoun et al. (2007), Mosquera et al. (2008), Landau et al. (2008), Landau & Scóccola (2010), Mosquera & Civitarese (2010), Civitarese et al. (2010), among others.

If the Higgs vacuum expectation value acquires a different value during BBN than the present one, the electron mass, the proton-neutron mass difference, the Fermi constant, and the deuterium binding energy, ϵ_D , will be different from the corresponding actual values. Several authors have studied the dependence of ϵ_D on v (Flambaum & Shuryak 2002, 2003; Dmitriev & Flambaum 2003; Beane & Savage 2003; Epelbaum et al. 2003; Yoo & Scherrer 2003; Dmitriev et al. 2004; Flambaum & Wiringa 2007; Berengut et al. 2010; Mosquera & Civitarese 2010; Civitarese et al. 2010).

In this work, we study the effects of a possible variation of the fine structure constant and of the Higgs vacuum expectation value, considering a fixed value of the strong-coupling constant Λ_{QCD} . To perform the calculation of the primordial abundances, we use the linear dependence of ϵ_D with v discussed in Mosquera & Civitarese (2010); Civitarese et al. (2010). We use observational data of D, ^4He , and ^7Li to obtain constraints on the variation of the participant fundamental constants. We also perform an analysis of the sensibility of these constraints on the lithium abundance and the dependencies between ϵ_D and v .

Table 1. Values of the coefficient κ of Eq. (1).

Potential	κ
Argonne	-1.23
Bonn	-0.66
Nijmegen	-1.66
Reid	-1.83

This work is organized as follows. In Sect. 2 we calculate the primordial abundances and obtain constraints on the joint variation of α and v . The conclusions are presented in Sect. 3.

2. Bounds from BBN

In this section we study the effect of a possible variation of the fine structure constant, α , and of the Higgs vacuum expectation value, v , during primordial nucleosynthesis. If α acquires a value different from the actual one, the neutron-to-proton ratio, the neutron-proton mass difference, the weak decay rates, and the cross sections of several reactions involved during BBN would be different from current estimates (Landau et al. 2008), and consequently, the primordial abundances would be modified. The electron mass, the Fermi constant, the neutron-proton mass difference, the weak decay rates, the masses of light nuclei, the deuterium binding energy, ϵ_D , and the initial abundance of deuterium will be modified if the Higgs vacuum expectation value varies with time (Landau et al. 2008; Mosquera & Civitaresse 2010).

The dependence of the deuterium-binding energy on the Higgs vacuum expectation value is model dependent (Flambaum & Wiringa 2007). In this work, we use the results obtained by Mosquera & Civitaresse (2010); Civitaresse et al. (2010). The authors had calculated the constant of proportionality, κ , which relates ϵ_D and v , for four different nucleon-nucleon potentials (Argonne v_{18} potential, Bonn potential, Nijmegen potential and Reid 93 potential),

$$\frac{\Delta\epsilon_D}{(\epsilon_D)_0} = \kappa \frac{\Delta v}{v_0}, \quad (1)$$

where $\Delta\epsilon_D = (\epsilon_D)_{\text{BBN}} - (\epsilon_D)_0$, and $\Delta v = v_{\text{BBN}} - v_0$. The subindexes BBN and 0 indicate the value of the constant at primordial nucleosynthesis and at the present time, respectively. In Table 1 we present the values of κ obtained by Mosquera & Civitaresse (2010); Civitaresse et al. (2010). The Argonne v_{18} potential includes an electromagnetic interaction (Wiringa et al. 1995), proportional to the fine structure constant. We modified this potential to include the variation of the fine structure constant and performed the calculation of ϵ_D . We obtained the relation $\frac{\Delta\epsilon_D}{(\epsilon_D)_0} = -0.0019 \frac{\Delta\alpha}{\alpha_0}$.

In order to calculate the primordial abundances, we modified the code developed by Kawano (Kawano 1988, 1992) for each nucleon-nucleon potential of Table 1 (Mosquera & Civitaresse 2010; Civitaresse et al. 2010). For details about the corrections introduced in the public version of Kawano's code see Mosquera et al. (2008); Landau et al. (2008); Mosquera & Civitaresse (2010).

The WMAP data are able to constraint the baryon density $\Omega_B h^2$ (related to the baryon-to-photon ratio η_B) with great accuracy, but there is still some degeneracy between the model parameters, namely: one is dealing with a parametric hypersurface defined by the values of $\Omega_B h^2$, $\Omega_{\text{CDM}} h^2$ (dark matter density in units of the critical density), Θ (gives the ratio of the comoving

Table 2. Best-fit parameter values for the BBN constraints.

κ	$\eta_B \pm \sigma [10^{-10}]$	$\frac{\Delta\alpha}{\alpha_0} \pm \sigma [10^{-3}]$	$\frac{\Delta v}{v_0} \pm \sigma [10^{-3}]$
-1.23	6.440 ^{+0.382} _{-0.219}	-2.5 ± 4.8	29.5 ^{+1.3} _{-1.1}
-0.66	6.150 ^{+0.365} _{-0.209}	-8.5 ^{+4.5} _{-4.7}	29.9 ^{+1.3} _{-1.1}
-1.66	6.744 ^{+0.317} _{-0.378}	1.5 ^{+4.7} _{-4.0}	29.6 ^{+1.1} _{-1.3}
-1.83	6.901 ^{+0.243} _{-0.386}	3.0 ± 4.0	29.6 ^{+1.0} _{-1.4}

sound horizon at decoupling to the angular diameter distance to the surface of last scattering), τ (reionization optical depth), n_s (scalar spectral index), A_s (amplitude of the density fluctuations). For this reason we computed the light nuclei abundances for the following cases:

- i) variation of α and v allowing η_B vary,
- ii) variation of α and v keeping η_B fixed at WMAP value ($\eta_B^{\text{WMAP}} = (6.108 \pm 0.219) \times 10^{-10}$) (Spergel et al. 2003, 2007).

In order to obtain the best-fit values for the parameters, we performed a χ^2 -test to compare the theoretical abundances and the observational data.

The observational data for D were extracted from Burles & Tytler (1998a,b), O'Meara et al. (2001), Pettini & Bowen (2001), Levshakov et al. (2002), Kirkman et al. (2003), Crighton et al. (2004), O'Meara et al. (2006), Pettini et al. (2008), Ivanchik et al. (2010).

We used the data from Izotov et al. (2006), Peimbert et al. (2007), Izotov et al. (2007), Izotov & Thuan (2010) for ^4He , and for ^7Li we considered the data given by Bonifacio & Molaro (1997), Molaro et al. (1997), Ryan et al. (2000), Bonifacio et al. (2002), Boesgaard et al. (2005), Asplund et al. (2006), Bonifacio et al. (2007), Hosford et al. (2009). Regarding the consistency of the data, we followed the treatment of Yao et al. (2006) and increased the errors by a fixed factor: $\Theta_D = 2.37$, $\Theta_{^4\text{He}} = 2.69$ and $\Theta_{^7\text{Li}} = 1.43$ for D, ^4He and ^7Li , respectively.

2.1. Variation of α and v allowing η_B to vary

We calculated the BBN abundances for different values of the fine structure constant, the Higgs vacuum expectation value and η_B , for each potential considered (see Table 1). We performed a χ^2 -test to find the best-fit values of the parameters. In Table 2 we show the results of the parameters corresponding to the best fit, at 1σ , for the BBN constraints on η_B (in units of 10^{-10}), $\frac{\Delta\alpha}{\alpha_0}$ (in units of 10^{-3}) and $\frac{\Delta v}{v_0}$ (in units of 10^{-3}). The values were obtained for the different values of κ given in Table 1. For all these cases, the value of $\frac{\chi^2_{\text{min}}}{N-3}$ is 1.00.

We found good agreement at the level of three standard deviations between our best-fit value of η_B and the one obtained using WMAP data (Spergel et al. 2007). We also found null variation of α within two standard deviations, and variation of the Higgs vacuum expectation value, even at the level of six standard deviations. In Fig. 1 we present the 3σ -likelihood contours for η_B , $\frac{\Delta\alpha}{\alpha_0}$ and $\frac{\Delta v}{v_0}$, and the one-dimensional likelihood for all potentials.

2.2. Variation of α and v keeping η_B fixed

Once again, we calculated the BBN abundances for different values of the fine structure constant and of the Higgs vacuum




Fig. 1. 3σ Likelihood contours for $\frac{\Delta\alpha}{\alpha_0}$, $\frac{\Delta v}{v_0}$ and η_B and one-dimensional likelihood for all values of κ .

Table 3. Best-fit parameters with η_B fixed by WMAP.

κ	$\frac{\Delta\alpha}{\alpha_0} \pm \sigma [10^{-3}]$	$\frac{\Delta v}{v_0} \pm \sigma [10^{-3}]$	$\frac{\chi^2_{\min}}{N-2}$
-1.23	-2.0 ± 4.0	29.0 ± 1.5	1.04
-0.66	-9.0 ± 5.0	29.5 ± 2.0	0.96
-1.66	3.0 ± 4.0	28.5 ± 1.5	1.19
-1.83	5.0 ± 4.0	28.0 ± 2.0	1.25

expectation value for each potential considered (see Table 1). This calculation was made keeping η_B fixed at WMAP value ($\eta_B^{\text{WMAP}} = (6.108 \pm 0.219) \times 10^{-10}$) (Spergel et al. 2007). We performed a χ^2 -test to find the best-fit value. In Table 3 we present the results for the best-fit parameter values within one standard deviation for the BBN constraints on $\frac{\Delta\alpha}{\alpha_0}$ (in units of 10^{-3}) and $\frac{\Delta v}{v_0}$ (in units of 10^{-3}), with η_B fixed at the WMAP estimate, for the different values of κ , as given in Table 1.

In Fig. 2 we present the 3σ -likelihood contours for $\frac{\Delta\alpha}{\alpha_0}$ and $\frac{\Delta v}{v_0}$ and one-dimensional likelihood, for all potentials.

We found null variation of α at 2σ level, but the variation of v is non-null even at 6σ . If the analysis does not include the lithium-data-set, we found null variation of both fundamental constants at the level of 2σ .

3. Conclusion

We obtained bounds on the joint variation of α and v using the observational abundances of D, ^4He , and ^7Li . We performed the analysis for different estimates of the dependence of the deuterium-binding energy on the Higgs vacuum expectation value. We found that the four dependencies considered, that is for the four values of κ of Table 1, give similar results. We found reasonable fits for the variation of α , v and η_B for the whole data set. However, and referring to the particular case of v , its variation is significant only when the data-set includes the ^7Li abundance. If the present values of ^7Li abundances are correct (Meléndez & Ramírez 2004; Richard et al. 2005; Prodanović & Fields 2007), then varying fundamental constants




Fig. 2. 3σ Likelihood contours for $\frac{\Delta\alpha}{\alpha_0}$ vs. $\frac{\Delta v}{v_0}$ and one-dimensional likelihood for all values of κ .

would be a possible explanation to solve the discrepancy between the light element abundances and the WMAP estimates.

Acknowledgements. Support for this work was provided by the PIP 0740 of the National Research Council (CONICET) of Argentina. The authors are members of the Scientific Research Career of the CONICET.

References

- Asplund, M., Lambert, D. L., Nissen, P. E., Primas, F., & Smith, V. V. 2006, *ApJ*, 644, 229
- Bahcall, J. N., Steinhardt, C. L., & Schlegel, D. 2004, *ApJ*, 600, 520
- Barr, S. M., & Mohapatra, P. K. 1988, *Phys. Rev. D*, 38, 3011
- Beane, S. R., & Savage, M. J. 2003, *Nucl. Phys. A*, 713, 148
- Berengut, J. C., Flambaum, V. V., & Dmitriev, V. F. 2010, *Phys. Lett. B*, 683, 114
- Bergström, L., Iguri, S., & Rubinstein, H. 1999, *Phys. Rev. D*, 60, 45005
- Bize, S., Diddams, S. A., Tanaka, U., et al. 2003, *Phys. Rev. Lett.*, 90, 150802
- Boesgaard, A. M., Novicki, M. C., & Stephens, A. 2005, in *From Lithium to Uranium: Elemental Tracers of Early Cosmic Evolution*, ed. V. Hill, P. François, & F. Primas, *IAU Symp.*, 228, 29
- Bonifacio, P., & Molaro, P. 1997, *MNRAS*, 285, 847
- Bonifacio, P., Pasquini, L., Spite, F., et al. 2002, *ApJ*, 390, 91
- Bonifacio, P., Molaro, P., Sivarani, T., et al. 2007, *A&A*, 462, 851
- Brax, P., van de Bruck, C., Davis, A.-C., & Rhodes, C. S. 2003, *Astro. Nucl. Space Sci.*, 283, 627
- Burles, S., & Tytler, D. 1998a, *ApJ*, 499, 699
- Burles, S., & Tytler, D. 1998b, *ApJ*, 507, 732
- Campbell, B. A., & Olive, K. A. 1995, *Phys. Lett. B*, 345, 429
- Chamoun, N., Landau, S. J., Mosquera, M. E., & Vucetich, H. 2007, *J. Phys. G Nucl. Phys.*, 34, 163
- Civitarese, O., Moliné, M. A., & Mosquera, M. E. 2010, *Nucl. Phys., A*, 846, 157
- Coc, A., Nunes, N. J., Olive, K. A., Uzan, J.-P., & Vangioni, E. 2007, *Phys. Rev. D*, 76, 023511
- Crighton, N. H. M., Webb, J. K., Ortiz-Gil, A., & Fernández-Soto, A. 2004, *MNRAS*, 355, 1042
- Cybur, R. H., Fields, B. D., Olive, K. A., & Skillman, E. 2005, *Astropart. Phys.*, 23, 313
- Damour, T., & Dyson, F. 1996, *Nucl. Phys. B*, 480, 37
- Damour, T., & Polyakov, A. M. 1994, *Nucl. Phys. B*, 95, 10347
- Damour, T., Piazza, F., & Veneziano, G. 2002a, *Phys. Rev. Lett.*, 89, 081601
- Damour, T., Piazza, F., & Veneziano, G. 2002b, *Phys. Rev. D*, 66, 046007
- Dmitriev, V. F., & Flambaum, V. V. 2003, *Phys. Rev. D*, 67, 063513
- Dmitriev, V. F., Flambaum, V. V., & Webb, J. K. 2004, *Phys. Rev. D*, 69, 063506
- Epelbaum, E., Meißner, U., & Glöckle, W. 2003, *Nucl. Phys. A*, 714, 535
- Fischer, M., Kolachevsky, N., Zimmermann, M., et al. 2004, *Phys. Rev. Lett.*, 92, 230802

- Flambaum, V. V., & Shuryak, E. V. 2002, *Phys. Rev. D*, 65, 103503
 Flambaum, V. V., & Shuryak, E. V. 2003, *Phys. Rev. D*, 67, 083507
 Flambaum, V. V., & Wiringa, R. B. 2007, *Phys. Rev. C*, 76, 054002
 Fujii, Y., Iwamoto, A., Fukahori, T., et al. 2000, *Nucl. Phys. B*, 573, 377
 Gleiser, M., & Taylor, J. G. 1985, *Phys. Rev. D*, 31, 1904
 Hosford, A., Ryan, S. G., García Pérez, A. E., Norris, J. E., & Olive, K. A. 2009, *A&A*, 493, 601
 Ichikawa, K., & Kawasaki, M. 2002, *Phys. Rev.*, D65, 123511
 Ichikawa, K., & Kawasaki, M. 2004, *Phys. Rev. D*, 69, 123506
 Ivanchik, A., Petitjean, P., Varshalovich, D., et al. 2005, *A&A*, 440, 45
 Ivanchik, A. V., Petitjean, P., Balashev, S. A., et al. 2010, *MNRAS, Soc.*, 297
 Izotov, Y. I., & Thuan, T. X. 2010, *ApJ*, 710, L67
 Izotov, Y. I., Schaerer, D., Blecha, A., et al. 2006, *A&A*, 459, 71
 Izotov, Y. I., Thuan, T. X., & Stasińska, G. 2007, *ApJ*, 662, 15
 Kaluza, T. 1921, *Sitzungber. Preuss. Akad. Wiss. K*, 1, 966
 Kawano, L. 1988, fERMILAB-PUB-88-034-A
 Kawano, L. 1992, fERMILAB-PUB-92-004-A
 King, J. A., Webb, J. K., Murphy, M. T., & Carswell, R. F. 2008, *Phys. Rev. Lett.*, 101, 251304
 Kirkman, D., Tytler, D., Suzuki, N., O'Meara, J. M., & Lubin, D. 2003, *ApJSS*, 149, 1
 Klein, O. 1926, *Z. Phys.*, 37, 895
 Landau, S. J., & Scóccola, G. 2010, *A&A*, 517, A62
 Landau, S. J., Mosquera, M. E., & Vucetich, H. 2006, *ApJ*, 637, 38
 Landau, S. J., Mosquera, M. E., Scóccola, C. G., & Vucetich, H. 2008, *Phys. Rev. D*, 78, 083527
 Levshakov, S. A., Dessauges-Zavadsky, M., D'Odorico, S., & Molaro, P. 2002, *ApJ*, 565, 696
 Müller, C. M., Schäfer, G., & Wetterich, C. 2004, *Phys. Rev. D*, 70, 083504
 Maeda, K. 1988, *Modern Phys. Lett. A*, 31, 243
 Malec, A. L., Buning, R., Murphy, M. T., et al. 2010, *MNRAS*, 403, 1541
 Marion, H., Pereira Dos Santos, F., Abgrall, M., et al. 2003, *Phys. Rev. Lett.*, 90, 150801
 Martínez Fiorenzano, A. F., Vladilo, G., & Bonifacio, P. 2003, *Soc. Astron. Ital. Mem. Suppl.*, 3, 252
 Meléndez, J., & Ramírez, I. 2004, *ApJ*, 615, L33
 Molaro, P., Bonifacio, P., & Pasquini, L. 1997, *MNRAS*, 292, L1
 Mosquera, M. E., & Civitaresse, O. 2010, *A&A*, 520, A112
 Mosquera, M. E., Scóccola, C., Landau, S., & Vucetich, H. 2008, *A&A*, 478, 675
 Murphy, M. T., Webb, J. K., Flambaum, V. V., et al. 2001a, *MNRAS*, 327, 1208
 Murphy, M. T., Webb, J. K., Flambaum, V. V., Prochaska, J. X., & Wolfe, A. M. 2001b, *MNRAS*, 327, 1237
 Murphy, M. T., Webb, J. K., & Flambaum, V. V. 2003, *MNRAS*, 345, 609
 Nollett, K. M., & Lopez, R. E. 2002, *Phys. Rev.*, D66, 063507
 O'Meara, J. M., Tytler, D., Kirkman, D., et al. 2001, *ApJ*, 552, 718
 O'Meara, J. M., Burles, S., Prochaska, J. X., et al. 2006, *ApJ*, 649, L61
 Overduin, J. M., & Wesson, P. S. 1997, *Phys. Rep.*, 283, 303
 Palma, G. A., Brax, P., Davis, A. C., & van de Bruck, C. 2003, *Phys. Rev. D*, 68, 123519
 Peik, E., Lipphardt, B., Schnatz, H., et al. 2004, *Phys. Rev. Lett.*, 93, 170801
 Peimbert, M., Luridiana, V., & Peimbert, A. 2007, *ApJ*, 666, 636
 Pettini, M., & Bowen, D. V. 2001, *ApJ*, 560, 41
 Pettini, M., Zych, B. J., Murphy, M. T., Lewis, A., & Steidel, C. C. 2008, *MNRAS*, 391, 1499
 Prestage, J. D., Tjoelker, R. L., & Maleki, L. 1995, *Phys. Rev. Lett.*, 74, 3511
 Prodanović, T., & Fields, B. D. 2007, *Phys. Rev. D*, 76, 083003
 Quast, R., Reimers, D., & Levshakov, S. A. 2004, *A&A*, 415, L7
 Reinhold, E., Buning, R., Hollenstein, U., et al. 2006, *Phys. Rev. Lett.*, 96, 151101
 Richard, O., Michaud, G., & Richer, J. 2005, *ApJ*, 619, 538
 Ryan, S. G., Beers, T. C., Olive, K. A., Fields, B. D., & Norris, J. E. 2000, *ApJ*, 530, L57
 Sortais, Y., Bize, S., Abgrall, M., et al. 2000, *Phys. Scripta*, T95, 50
 Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJSS*, 148, 175
 Spergel, D. N., Bean, R., Doré, O., et al. 2007, *ApJSS*, 170, 377
 Srianand, R., Chand, H., Petitjean, P., & Aracil, B. 2004, *Phys. Rev. Lett.*, 92, 121302
 Thompson, R. I., Bechtold, J., Black, J. H., et al. 2009, *ApJ*, 703, 1648
 Tzanavaris, P., Murphy, M. T., Webb, J. K., Flambaum, V. V., & Curran, S. J. 2007, *MNRAS*, 374, 634
 Webb, J. K., Flambaum, V. V., Churchill, C. W., Drinkwater, M. J., & Barrow, J. D. 1999, *Phys. Rev. Lett.*, 82, 884
 Webb, J. K., Murphy, M. T., Flambaum, V. V., et al. 2001, *Phys. Rev. Lett.*, 87, 091301
 Weinberg, S. 1983, *Phys. Lett. B*, 125, 265
 Wiringa, R. B., Stoks, V. G. J., & Schiavilla, R. 1995, *Phys. Rev. C*, 51, 38
 Wu, Y., & Wang, Z. 1986, *Phys. Rev. Lett.*, 57, 1978
 Yao, W.-M., Amsler, C., Asner, D., et al. 2006, *J. Phys. G*, 33, 1
 Yoo, J. J., & Scherrer, R. J. 2003, *Phys. Rev. D*, 67, 043517
 Youm, D. 2001a, *Phys. Rev. D*, 63, 125011
 Youm, D. 2001b, *Phys. Rev. D*, 64, 085011